

# TOWARDS COMMUTATOR THEORY FOR RELATIONS. IV.

PAOLO LIPPARINI

**ABSTRACT.** We provide more characterizations of varieties having a term Mal'cev modulo two functions  $F$  and  $G$ . We characterize varieties neutral in the sense of  $F$ , that is varieties satisfying  $R \subseteq F(R)$ . We present examples of global operators satisfying the homomorphism property, in particular we show that many known commutators satisfy the homomorphism property.

See Parts I-III [2] for unexplained notation.

We now find more conditions equivalent to the existence of a term Mal'cev modulo  $F$  and  $G$  throughout a variety, thus complementing Theorem 3 in Part III.

**Theorem 1.** *Suppose that  $\mathcal{V}$  is a variety,  $F, G$  are global operators on  $\mathcal{V}$  for admissible and reflexive relations,  $F, G$  are monotone and satisfy the homomorphism property. Then the following are equivalent:*

(i)  $\mathcal{V}$  has a term which is Mal'cev modulo  $F_{\mathbf{A}}$  and  $G_{\mathbf{A}}$  for every algebra  $\mathbf{A}$  in  $\mathcal{V}$ .

(ii) In every algebra  $\mathbf{A} \in \mathcal{V}$  and for all relations  $R, S, T \in \text{Adm}(\mathbf{A})$ , the following holds:

$$T(R \circ S) \subseteq F_{\mathbf{A}}(T) \circ \overline{S^- \cup R} \circ G_{\mathbf{A}}(S).$$

(iii) In the free algebra  $\mathbf{X}$  in  $\mathcal{V}$  generated by 2 elements the following holds:

$$T(R \circ S) \subseteq F_{\mathbf{X}}(T) \circ S^- \circ R \circ G_{\mathbf{X}}(S),$$

for all relations  $R, S, T \in \text{Adm}(\mathbf{X})$ .

(iv) In every algebra  $\mathbf{A} \in \mathcal{V}$  and for all relations  $S, T \in \text{Adm}(\mathbf{A})$ , the following holds:

$$TS \subseteq F_{\mathbf{A}}(T) \circ S^- \circ G_{\mathbf{A}}(T).$$

(v) In the free algebra  $\mathbf{X}$  in  $\mathcal{V}$  generated by 2 elements the following holds:

$$TS \subseteq F_{\mathbf{X}}(T) \circ S^- \circ G_{\mathbf{X}}(T),$$

for all relations  $S, T \in \text{Adm}(\mathbf{X})$ .

---

2000 *Mathematics Subject Classification.* Primary 08A99; 08B99.

*Key words and phrases.* Neutral varieties, weak difference term, Mal'cev term, global operator satisfying the homomorphism property, commutators.

The author has received support from MPI and GNSAGA.

(vi) In the free algebra  $\mathbf{X}$  in  $\mathcal{V}$  generated by the 3 elements  $x, y, z$  the following holds:

$$S \circ T \subseteq F_{\mathbf{X}}(S) \circ Cg(T) \circ Cg(S) \circ G_{\mathbf{X}}(T).$$

where  $S$  is the smallest reflexive and admissible relation on  $\mathbf{X}$  containing  $(x, y)$ , and  $T$  is the smallest reflexive and admissible relation on  $\mathbf{X}$  containing  $(y, z)$ .

(vii) In the free algebra  $\mathbf{X}$  in  $\mathcal{V}$  generated by the 3 elements  $x, y, z$  the following holds:

$$S \circ S \subseteq F_{\mathbf{X}}(S) \circ \Theta \circ G_{\mathbf{X}}(S).$$

where  $S$  is the smallest reflexive and admissible relation on  $\mathbf{X}$  containing  $(x, y)$  and  $(y, z)$ , and  $\Theta$  is the tolerance generated by  $S$ .

*Proof.* (i)  $\Rightarrow$  (ii). If  $(a, c) \in T(R \circ S)$  then  $a T c$  and there exists  $b \in A$  such that  $a R b S c$ . Thus,  $a F(T) t(a, c, c) \overline{S^- \cup R} t(b, b, c) G(S) c$ . Actually, (i)  $\Rightarrow$  (ii) is a particular case of Theorem 1(i) in Part III.

(ii)  $\Rightarrow$  (iii) and (iv)  $\Rightarrow$  (v) are trivial.

Both (ii)  $\Rightarrow$  (iv) and (iii)  $\Rightarrow$  (v) are obtained by taking  $R = 0$ .

(v)  $\Rightarrow$  (i). Let  $x, y$  be the generators of  $\mathbf{X}$ . By taking, in (v),  $T = S$  to be the smallest admissible and reflexive relation of  $\mathbf{X}$  containing  $(x, y)$  we get Condition (x) in Part III, Theorem 3, which implies (i) by the same theorem.

(i)  $\Rightarrow$  (vi) and (i)  $\Rightarrow$  (vii) are immediate from Part III, Theorem 1 (iii).

(vi)  $\Rightarrow$  (i). Since  $x S y T z$ , we have  $(x, z) \in S \circ T$ , hence  $(x, z) \in F_{\mathbf{X}}(S) \circ Cg(T) \circ Cg(S) \circ G_{\mathbf{X}}(T)$  by assumption.

This means that there are terms  $t_1(x, y, z)$ ,  $t(x, y, z)$  and  $t_2(x, y, z)$  such that  $(x, t_1(x, y, z)) \in F_{\mathbf{X}}(S)$ ,  $(t_1(x, y, z), t(x, y, z)) \in Cg(T)$ ,  $(t(x, y, z), t_2(x, y, z)) \in Cg(S)$  and  $(t_2(x, y, z), z) \in G_{\mathbf{X}}(T)$ .

Since  $Cg(T) = \{(u(x, y, z), v(x, y, z)) | u, v \text{ terms of } \mathcal{V} \text{ such that } u(x, y, y) = v(x, y, y)\}$ , we get  $t_1(x, y, y) = t(x, y, y)$ , and similarly  $t(y, y, z) = t_2(y, y, z)$ .

Now the proof goes exactly as in the final part of the proof of Part III, Theorem 3 (vii)  $\Rightarrow$  (i).

(vii)  $\Rightarrow$  (i). Let  $\mathbf{X}, S, \Theta$  be as in (vii). Since  $x S y S z$ , we have  $(x, z) \in S \circ S$ , hence  $(x, z) \in F_{\mathbf{X}}(S) \circ \Theta \circ G_{\mathbf{X}}(S)$ .

This means that  $\mathcal{V}$  has terms  $t_1(x, y, z)$  and  $t_2(x, y, z)$  such that  $(x, t_1(x, y, z)) \in F_{\mathbf{X}}(S)$ ,  $(t_1(x, y, z), t_2(x, y, z)) \in \Theta$  and  $(t_2(x, y, z), z) \in G_{\mathbf{X}}(S)$ .

Notice that  $\Theta = \{(u(x, y, z, x, y, y, z), u(x, y, z, y, x, z, y)) | u \text{ a term of } \mathbf{X}\}$ . Indeed, the right-hand relation is reflexive, symmetric, admissible, and contains  $(x, y)$  and  $(y, z)$ ; moreover, every other reflexive, symmetric admissible relation containing  $(x, y)$  and  $(y, z)$  contains all pairs of the form  $(u(x, y, z, x, y, y, z), u(x, y, z, y, x, z, y))$ .

Hence,  $(t_1(x, y, z), t_2(x, y, z)) \in \Theta$  means that there is a 7-ary term  $t'$  such that  $t_1(x, y, z) = t'(x, y, z, x, y, y, z)$  and  $t'(x, y, z, y, x, z, y) = t_2(x, y, z)$ .

We claim that the ternary term  $t(x, y, z) = t'(x, y, z, x, y, z, y)$  is Mal'cev modulo  $F$  and  $G$  throughout  $\mathcal{V}$ . First, notice that  $t_1(x, y, y) = t(x, y, y)$ , and

$t(y, y, z) = t_2(y, y, z)$ . Now the last part of the proof of Part III, Theorem 3 (vii)  $\Rightarrow$  (i) shows that  $a F_{\mathbf{A}}(R) t_1(a, b, b) = t(a, b, b)$  holds for every algebra  $\mathbf{A} \in \mathcal{V}$  and  $a R b$ , and a similar relation holds for  $G$ , thus  $t$  is Mal'cev modulo  $F$  and  $G$ .  $\square$

We say that  $F$  is *regular* on  $\mathcal{V}$  if and only if  $F$  is a global operator on  $\mathcal{V}$  from admissible and reflexive relations to congruences (that is,  $F_{\mathbf{A}}(R)$  is always supposed to be a congruence of  $\mathbf{A}$ ),  $F$  is monotone, satisfies the homomorphism property, and satisfies also the following *quotient property*: “for every  $\mathbf{A} \in \mathcal{V}$  and every  $R \in \text{Adm}(\mathbf{A})$ , it happens that  $F_{\mathbf{B}}(R/F_{\mathbf{A}}(R)) = 0_{\mathbf{B}}$ , where  $\mathbf{B}$  is  $\mathbf{A}/F_{\mathbf{A}}(R)$ ”. Here,  $R/F_{\mathbf{A}}(R)$  denotes  $\pi(R)$ , where  $\pi$  denotes the canonical epimorphism  $\mathbf{A} \rightarrow \mathbf{A}/F_{\mathbf{A}}(R)$ .

In the particular case when  $F = G$ , we simply say that a term  $t$  is *Mal'cev modulo  $F$*  in order to mean that  $t$  is Mal'cev modulo  $F$  and  $F$ .

**Theorem 2.** *Suppose that  $\mathcal{V}$  is a variety and  $F$  is a global operator which is regular on  $\mathcal{V}$  and satisfies  $F(R) = F(R^-)$ , for every admissible relation  $R$ . Then the following are equivalent:*

- (i)  $\mathcal{V}$  has a term which is Mal'cev modulo  $F_{\mathbf{A}}$  for every algebra  $\mathbf{A}$  in  $\mathcal{V}$ .
- (ii) For every  $\mathbf{A} \in \mathcal{V}$  and  $R \in \text{Adm}(\mathbf{A})$ , if  $F_{\mathbf{A}}(R) = 0_{\mathbf{A}}$  then  $R$  is a congruence of  $\mathbf{A}$ .
- (iii) For every  $\mathbf{A} \in \mathcal{V}$  and  $R \in \text{Adm}(\mathbf{A})$ , if  $F_{\mathbf{A}}(R) = 0_{\mathbf{A}}$  then  $R$  is a tolerance of  $\mathbf{A}$ .
- (iv) For every  $\mathbf{A} \in \mathcal{V}$  and  $R \in \text{Adm}(\mathbf{A})$ , if  $F_{\mathbf{A}}(R) = 0_{\mathbf{A}}$  then  $R$  is a transitive relation of  $\mathbf{A}$ .
- (v) Let  $\mathbf{X}$  be the free algebra in  $\mathcal{V}$  generated by  $\{x, y\}$  and  $S$  be the smallest reflexive and admissible relation on  $\mathbf{X}$  containing  $(x, y)$ . If  $\mathbf{B} = \mathbf{X}/F_{\mathbf{X}}(S)$ , then  $T = S/F_{\mathbf{X}}(S)$  is (a) a congruence (equivalently, (b) a tolerance) of  $\mathbf{B}$ .
- (vi) Let  $\mathbf{X}$  be the free algebra in  $\mathcal{V}$  generated by  $\{x, y, z\}$  and  $S$  be the smallest reflexive and admissible relation on  $\mathbf{X}$  containing  $(x, y)$  and  $(y, z)$ . If  $\mathbf{B} = \mathbf{X}/F_{\mathbf{X}}(S)$ , then  $T = S/F_{\mathbf{X}}(S)$  is a transitive relation of  $\mathbf{B}$ .

*Proof.* (i)  $\Rightarrow$  (ii) is immediate from Part III, Theorem 1(xi).

(ii)  $\Rightarrow$  (iii) and (ii)  $\Rightarrow$  (iv) are trivial.

Let  $\mathbf{X}, \mathbf{B}, S, T$  be as in (v) (respectively, as in (vi)). By the quotient property,  $F_{\mathbf{B}}(T) = 0_{\mathbf{B}}$ , thus (ii), (iii), (iv) imply, respectively, (v)(a), (v)(b), (vi).

If  $T$  is a congruence then  $T$  is a tolerance, thus (v)(a) implies (v)(b).

Suppose that (v)(b) holds. Since  $T = S/F_{\mathbf{X}}(S)$  is a tolerance, and  $\pi$  is surjective, then  $\pi^{-1}(T) = \pi^{-1}(S/F_{\mathbf{X}}(S)) = F_{\mathbf{X}}(S) \circ S \circ F_{\mathbf{X}}(S)$  is a tolerance, hence  $S \subseteq F_{\mathbf{X}}(S) \circ S \circ F_{\mathbf{X}}(S) = (F_{\mathbf{X}}(S) \circ S \circ F_{\mathbf{X}}(S))^- = F_{\mathbf{X}}(S) \circ S^- \circ F_{\mathbf{X}}(S)$ .

Thus  $S \subseteq F_{\mathbf{X}}(S) \circ S^- \circ F_{\mathbf{X}}(S)$ , which is Condition (x) in Part III, Theorem 3, which implies (i).

Suppose that (vi) holds, that is,  $T$  is transitive. Then  $\pi^{-1}(T)$  is transitive, hence  $S \circ S \subseteq \pi^{-1}(T) \circ \pi^{-1}(T) = \pi^{-1}(T) = F_{\mathbf{X}}(S) \circ S \circ F_{\mathbf{X}}(S)$ .

This is Condition (vii) in Part III, Theorem 3, which implies (i).  $\square$

The next theorem provides a version of our results for what might be called *neutral* varieties, that is, varieties satisfying  $R \subseteq F(R)$  (notice that we are not necessarily assuming  $F(R) \subseteq R$ ).

**Theorem 3.** *Suppose that  $\mathcal{V}$  is a variety,  $F$  is a global operator on  $\mathcal{V}$  for admissible and reflexive relations, and  $F$  is monotone and satisfies the homomorphism property. Then the following are equivalent:*

(i)  $R \subseteq F_{\mathbf{A}}(R)$  holds for every algebra  $\mathbf{A}$  in  $\mathcal{V}$  and for every relation  $R \in \text{Adm}(\mathbf{A})$ .

(ii)  $R \subseteq F_{\mathbf{X}}(R)$  holds for every admissible and reflexive relation  $R$  in the free algebra  $\mathbf{X}$  in  $\mathcal{V}$  generated by 2 elements.

(iii)  $S \subseteq F_{\mathbf{X}}(S)$  holds in the free algebra  $\mathbf{X}$  in  $\mathcal{V}$  generated by the two elements  $x, y$ , where  $S$  is the smallest admissible and reflexive relation of  $\mathbf{X}$  containing  $(x, y)$ .

(iii)' In the free algebra  $\mathbf{X}$  in  $\mathcal{V}$  generated by the two elements  $x, y$ , if  $S$  is the smallest admissible and reflexive relation of  $\mathbf{X}$  containing  $(x, y)$  then  $(x, y) \in F_{\mathbf{X}}(S)$ .

In case  $F$  is regular, the preceding conditions are also equivalent to the following ones:

(iv) For every algebra  $\mathbf{A}$  in  $\mathcal{V}$  and for every relation  $R \in \text{Adm}(\mathbf{A})$ , if  $R \neq 0$  then  $F_{\mathbf{A}}(R) \neq 0$ .

(v) Let  $\mathbf{X}$  be the free algebra in  $\mathcal{V}$  generated by the two elements  $x, y$ , and let  $S$  be the smallest admissible and reflexive relation of  $\mathbf{X}$  containing  $(x, y)$ . Then  $S/F_{\mathbf{X}}(S) = 0$  in  $\mathbf{B} = \mathbf{X}/F_{\mathbf{X}}(S)$ .

*Proof.* (i)  $\Rightarrow$  (ii)  $\Rightarrow$  (iii)  $\Rightarrow$  (iii)' are trivial.

(iii)'  $\Rightarrow$  (i). Suppose that  $\mathbf{A} \in \mathcal{V}$ ,  $a, b \in A$ ,  $R \in \text{Adm}(\mathbf{A})$ , and  $a R b$ .

Since  $\mathbf{X}$  is the free algebra in  $\mathcal{V}$  generated by  $\{x, y\}$ , there is a homomorphism  $\phi : \mathbf{X} \rightarrow \mathbf{A}$  such that  $\phi(x) = a$  and  $\phi(y) = b$ .

Since, by (iii)',  $x F_{\mathbf{X}}(S) y$ , we have that  $\phi(x) \phi(F_{\mathbf{X}}(S)) \phi(y)$ , hence, by the homomorphism property,  $a F_{\mathbf{A}}(\phi(S)) b$ .

Since  $S$  is the admissible and reflexive relation generated by  $(x, y)$ , and  $\phi$  is a homomorphism, then  $\phi(S)$  is the admissible and reflexive relation generated by  $(\phi(x), \phi(y)) = (a, b)$ , and, since  $a R b$ , we have that  $\phi(S) \subseteq R$ ; thus, by the monotonicity of  $F_{\mathbf{A}}$ , we get  $F_{\mathbf{A}}(\phi(S)) \subseteq F_{\mathbf{A}}(R)$  and, eventually,  $a F_{\mathbf{A}}(R) b$ .

(i)  $\Rightarrow$  (iv) is trivial.

(iv)  $\Rightarrow$  (v). By the quotient property,  $F_{\mathbf{B}}(S/F_{\mathbf{X}}(S)) = 0_{\mathbf{B}}$ , thus  $S/F_{\mathbf{X}}(S) = 0_{\mathbf{B}}$  by (iv).

(v)  $\Rightarrow$  (iii).  $S/F_{\mathbf{X}}(S) = 0_{\mathbf{B}}$  in  $\mathbf{B}$  means  $S \subseteq F_{\mathbf{X}}(S)$  in  $\mathbf{X}$ .  $\square$

We now show that the commutators defined in [1] and in Part I satisfy the homomorphism property. Actually, we shall deal with  $n$ -ary global operators on  $\mathcal{V}$ , that is, operators depending on  $n$ -variables (as above, each variable is intended to be an admissible and reflexive relation). In this general situation, the *homomorphism property* reads  $\phi(F_{\mathbf{B}}(R_1, \dots, R_n)) \subseteq F_{\mathbf{A}}(\phi(R_1), \dots, \phi(R_n))$ , for every  $\mathbf{A}, \mathbf{B} \in \mathcal{V}$ , homomorphism  $\phi : \mathbf{B} \rightarrow \mathbf{A}$ , and

$R_1, \dots, R_n \in \text{Adm}(\mathbf{B})$ . Moreover,  $F$  is *monotone* if and only if  $R_1 \subseteq S_1, \dots, R_n \subseteq S_n$  imply  $F(R_1, \dots, R_n) \subseteq F(S_1, \dots, S_n)$ .

If  $\mathbf{A}$  is any algebra, and  $R, S$  are compatible and reflexive relations,  $M(R, S)$  is defined to be the set of all *matrices* of the form

$$\begin{vmatrix} t(\bar{a}, \bar{b}) & t(\bar{a}, \bar{b}') \\ t(\bar{a}', \bar{b}) & t(\bar{a}', \bar{b}') \end{vmatrix},$$

where  $\bar{a}, \bar{a}' \in A^h$ ,  $\bar{b}, \bar{b}' \in A^k$ , for some  $h, k \geq 0$ ,  $t$  is an  $h + k$ -ary term operation of  $\mathbf{A}$ , and  $\bar{a} R \bar{a}', \bar{b} S \bar{b}'$ . Here,  $\bar{a} R \bar{a}'$  means  $a_1 R a'_1, a_2 R a'_2, \dots$ .

For  $R, S, T, U$  compatible and reflexive relations, let

$$K(R, S; U; T) = \left\{ (z, w) \mid \begin{vmatrix} x & y \\ z & w \end{vmatrix} \in M(R, S), x U z, x T y, \right\}.$$

$K(R, S; 1; T)$  is denoted by  $K(R, S; T)$  in Part I, and various commutators for relations are constructed there from  $K(R, S; T)$ .

**Proposition 4.** *For every variety  $\mathcal{V}$ ,  $K(R, S; U; T)$  and  $K(R, S; T)$  are monotone global operators on  $\mathcal{V}$  satisfying the homomorphism property.*

*Proof.* We shall give the proof for  $K(R, S; U; T)$ . The proof for  $K(R, S; T)$  is similar and simpler (and, anyway, is the particular case  $U = 1$  constantly). The operators are trivially monotone.

We now prove that  $K(R, S; U; T)$  satisfies the homomorphism property. Suppose that  $\mathbf{A}, \mathbf{B} \in \mathcal{V}$ ,  $\phi : \mathbf{B} \rightarrow \mathbf{A}$ ,  $R, S, T, U \in \text{Adm}(\mathbf{B})$ ,  $x', w' \in \mathbf{A}$ , and  $(x', w') \in \phi(K(R, S; U; T))$ . We have to show that  $(x', w') \in K(\phi(R), \phi(S); \phi(U); \phi(T))$ .

That  $(x', w') \in \phi(K(R, S; U; T))$  means that  $\mathbf{B}$  has couples  $(x_1, w_1), \dots, (x_m, w_m) \in K(R, S; U; T)$  such that, in  $\mathbf{A}$ ,  $(x', w')$  belongs to the admissible and reflexive relation generated by  $(\phi(x_1), \phi(w_1)), \dots, (\phi(x_m), \phi(w_m))$ . This means that  $\mathbf{A}$  has an  $m$ -ary polynomial  $p$  such that  $x' = p(\phi(x_1), \dots, \phi(x_m))$ , and  $w' = p(\phi(w_1), \dots, \phi(w_m))$ .

Since  $(x_i, w_i) \in K(R, S; U; T)$  for every  $i = 1, \dots, m$ , then, by the definition of  $K(R, S; U; T)$ , there are matrices

$$\begin{vmatrix} t_i(\bar{a}_i, \bar{b}_i) & t_i(\bar{a}_i, \bar{b}'_i) \\ t_i(\bar{a}'_i, \bar{b}_i) & t_i(\bar{a}'_i, \bar{b}'_i) \end{vmatrix}$$

such that  $\bar{a}_i R \bar{a}'_i$ ,  $\bar{b}_i S \bar{b}'_i$ ,  $t_i(\bar{a}_i, \bar{b}_i) T t_i(\bar{a}_i, \bar{b}'_i)$ ,  $t_i(\bar{a}_i, \bar{b}_i) U t_i(\bar{a}'_i, \bar{b}_i)$ , and  $x_i = t_i(\bar{a}'_i, \bar{b}_i)$ ,  $w_i = t_i(\bar{a}'_i, \bar{b}'_i)$ .

For sake of brevity, let us write  $t_i(\phi \bar{a}_i, \phi \bar{b}_i)$  in place of  $t_i(\phi(a_{i1}), \phi(a_{i2}), \dots, \phi(b_{i1}), \phi(b_{i2}), \dots)$ . The matrix

$$\begin{vmatrix} p(t_1(\phi \bar{a}_1, \phi \bar{b}_1), \dots, t_m(\phi \bar{a}_m, \phi \bar{b}_m)) & p(t_1(\phi \bar{a}_1, \phi \bar{b}'_1), \dots, t_m(\phi \bar{a}_m, \phi \bar{b}'_m)) \\ p(t_1(\phi \bar{a}'_1, \phi \bar{b}_1), \dots, t_m(\phi \bar{a}'_m, \phi \bar{b}_m)) & p(t_1(\phi \bar{a}'_1, \phi \bar{b}'_1), \dots, t_m(\phi \bar{a}'_m, \phi \bar{b}'_m)) \end{vmatrix}$$

belongs to  $M(\phi(R), \phi(S))$ ; moreover, since  $\phi$  is a homomorphism, the above matrix equals

$$\begin{vmatrix} p(\phi(t_1(\bar{a}_1, \bar{b}_1)), \dots, \phi(t_m(\bar{a}_m, \bar{b}_m))) & p(\phi(t_1(\bar{a}_1, \bar{b}'_1)), \dots, \phi(t_m(\bar{a}_m, \bar{b}'_m))) \\ p(\phi(t_1(\bar{a}'_1, \bar{b}_1)), \dots, \phi(t_m(\bar{a}'_m, \bar{b}_m))) & p(\phi(t_1(\bar{a}'_1, \bar{b}'_1)), \dots, \phi(t_m(\bar{a}'_m, \bar{b}'_m))) \end{vmatrix}$$

Since  $t_i(\bar{a}_i, \bar{b}_i) T t_i(\bar{a}_i, \bar{b}_i')$ , and since  $T$  is admissible, the elements in the upper row of the matrix are  $\phi(T)$ -related; similarly, the elements in the left-hand column are  $\phi(U)$ -related. Since  $x_i = t_i(\bar{a}_i', \bar{b}_i)$ , and  $w_i = t_i(\bar{a}_i', \bar{b}_i')$ , then the second row of the matrix consists of the elements  $p(\phi(x_1), \dots, \phi(x_m)) = x'$ , and  $p(\phi(w_1), \dots, \phi(w_m)) = w'$ . This means that the matrix witnesses that  $(x', w') \in K(\phi(R), \phi(S); \phi(U); \phi(T))$ .  $\square$

**Proposition 5.** *If  $\mathcal{V}$  is a variety,  $F_1, F_2, \dots$  are  $n$ -ary global operators on  $\mathcal{V}$  for admissible and reflexive relations, and  $F_1, F_2, \dots$  satisfy the homomorphism property, then so do the following operators:*

- (i)  $G_1(\bar{R}) = F_1(\bar{R}) \circ F_2(\bar{R})$ ;
- (ii)  $G_2(\bar{R}) = F_1(\bar{R}) \cap F_2(\bar{R})$ ;
- (iii)  $G_3(\bar{R}) = F_1(\bar{R}) \cup F_2(\bar{R})$ ;
- (iv)  $G_4(\bar{R}) = F_1(\bar{R}) + F_2(\bar{R})$ ;
- (v)  $G_5(\bar{R}) = (F_1(\bar{R}))^*$ ;
- (vi)  $G_6(\bar{R}) = K(F_1(\bar{R}), F_2(\bar{R}); F_3(\bar{R}); F_4(\bar{R}))$ .
- (vii)  $G_7(\bar{R}) = K(F_1(\bar{R}), F_2(\bar{R}); F_4(\bar{R}))$ .

*If, in addition,  $F_1, F_2, \dots$  are monotone, then so are  $G_1, \dots, G_7$ .*

*Proof.* The proof is similar to the proof of Proposition 4, in many cases simpler.  $\square$

Since, trivially, the constant operators  $F(\bar{R}) = 0$  and  $F(\bar{R}) = 1$ , the operator  $F(R) = R^-$ , as well as the projections, are monotone and satisfy the homomorphism property, we get from Proposition 5(vii)(v) that, for example,  $F(R, S) = K(R, S; 0)^*$  is monotone and satisfies the homomorphism property. Similarly, Proposition 5 can be iterated in order to show that all the commutators introduced in Part I are monotone and satisfy the homomorphism property. Notice also that condition (vi) in Proposition 5 is more general than Proposition 4: just take  $n = 4$ , and let each  $F_i$  be the projection onto the  $i$ -th coordinate.

## REFERENCES

- [1] Paolo Lipparini, *Congruence identities satisfied in  $n$ -permutable varieties*, Bollettino U. M. I. (7) **8-B** (1994), 851-868.
- [2] ———, *Towards commutator theory for relations*, <http://arxiv.org/math.GM/0507039>; II, <http://arxiv.org/math.GM/0509233>; III, <http://arxiv.org/math.GM/0511313> (2005).
- [3] ———, *Terms which are Mal'cev modulo some function*, submitted.

DIPARTIMENTO DI MATEMATICA, VIALE DELLA RICERCA SCIENTIFICA, II UNIVERSITÀ DI ROMA (OR VERGATTA), ROME ITALY

URL: <http://www.mat.uniroma2.it/~lipparin>